

In This Issue

Heel Pain After Long Winter

Keep Kids' Feet & Ankles Safe

Foot Fractures & Osteoporosis

Watch for Heel Pain After Coming Out of Hibernation

After a long winter, nothing beats going outside for a walk or run to enjoy the warm spring weather. However, sudden increased activity following several months of low or no activity can result in heel pain, also known as plantar fasciitis.


This painful condition results from inflammation of the tissue band (the plantar fascia) that extends from the heel to the toes. Repetitive activities, such as a new exercise routine or walking on a daily basis, can put stress and strain on the ligaments in the foot, leading to inflammation and pain.

The good news is heel pain can usually be relieved using conservative methods, but it must be treated early.

Heel pain can become chronic and debilitating if not cared for properly.

Our office can help you find relief with therapies, such as:

- anti-inflammatory medications
- stretching exercises
- orthotic devices
- physical therapy
- footwear modifications
- activity limitations



Although most patients with plantar fasciitis respond to nonsurgical treatment, some require surgery. If you continue to have heel pain with nonsurgical treatment, we can discuss your surgical options.

Heel pain should not stop you from taking in the beauty of spring. Make an appointment with our office if you are experiencing heel pain so we can help you resume a healthy and active lifestyle.

Keep Kids' Feet & Ankles Safe During Spring Sports

Competitive youth sports often require many athletes to transition from winter activities to spring activities without considering the increased risk of incurring a foot or ankle injury. Moving from indoor to outdoor playing surfaces with varying impact can stress a young athlete's feet and ankles, and going from sport to sport without allowing time for the muscles and bones to rest can cause overuse injuries.

If your child plans to participate in a sport this spring after playing through the winter sports season, follow these six tips:

- 1. Get a preseason health and wellness checkup.** A medical evaluation before the season starts can help identify any health concerns that could possibly lead to injury.
- 2. Take it slow.** Ask the coach to gradually increase children's playing time during practice and to avoid pushing them full throttle. Your child's feet and ankles need to become accustomed to the activity level required for the sport.
- 3. Wear proper, broken-in shoes.** Different sports require different shoe gear. Wearing the appropriate, well-fitting, broken-in athletic shoes can eliminate heel and toe discomfort.

4. Check your child's technique. Watch for any changes in your child's form or technique. Ask the coach to notify you if your child is placing more weight on one side of the body or is limping.

5. Insist on open communication if your child has pain. Express to your child athlete that s/he should inform you and the coach of any pain or discomfort as soon as it occurs. Overuse injuries, such as Achilles tendonitis and shin splints, can be subtle and can develop over time.

6. If an injury occurs, remember RICE. An injured foot or ankle can often be healed with rest, ice, compression and elevation (RICE). If your child complains of foot or ankle pain, s/he should take a break from playing and allow time for recovery.



Foot Fractures Can Signal Osteoporosis



Unexplained foot pain or foot fracture can actually be an early sign of osteoporosis, a bone-thinning disease that weakens bones and can lead to fractures or breaks.

Osteoporosis often progresses without any symptoms or is not diagnosed until a person experiences pain from a bone fracture. Bones with osteoporosis are in a weakened state, and normal weightbearing actions, such as walking, can cause the bones in the foot to break. Many of the patients who come to our office with foot pain find out they actually have a hairline break in the bone (stress fracture), without having experienced an injury.


While osteoporosis is most commonly seen in women over age 50, younger people and men are also affected. Early symptoms

can include increased pain with walking accompanied by redness and swelling on the top of the foot.


If you are living with osteoporosis, it is important to protect your feet from stress fractures. Our office recommends the following to keep your feet safe and comfortable:

- 1. Wear shoes designed for the exercise or sport.** Athletic shoes that support your arch and cushion the heel are the best to wear. Custom orthotics may also help provide extra support and shock absorption.
- 2. Start new workouts gradually.** Be sure to stretch or warm up for your workout and start new exercise routines gradually. Increase your exercise intensity by only 10 percent each week. If your feet get sore, use rest, ice, compression and elevation (RICE).
- 3. Protect your feet from bacteria.** Sweaty shoes and public showers at the gym are breeding grounds for bacteria, including resistant strains like MRSA. Never go barefoot while in public areas, and be sure to cover cuts and cracks in the skin or ingrown toenails to help keep out germs. If you have a cut or scrape that becomes infected and does not heal in a timely manner, call our office to have it examined.

If you are suffering from unexplained foot pain, schedule an appointment with our office for a proper diagnosis.



**Martin Foot
and Ankle**



2300 Pleasant Valley Rd.
York, PA 17402

2069 Carlisle Rd.
York, PA 17408

1010 Eichelberger St.
Suite 4
Hanover, PA 17331

2112 Harrisburg Pike
Suite 321
Lancaster, PA 17601

1042 Lititz Pike
Lititz, PA 17543

717-757-3537 all locations
800-456-0076

www.martinfootandankle.com

